

A Medical Corporation

John M. Adams, Jr., M.D. Kevin M. Ehrhart, M.D. Michael B. Gerhardt, M.D. Ronald J. Gowey, M.D. Kenton S. Horacek, M.D. Thomas P. Knapp, M.D. T. Michael Lain, M.D. Bert R. Mandelbaum, M.D. M. Ramin Modabber, M.D. John R. Sellman, M.D.

POST-OPERATIVE PROTOCOL ACL RECONSTRUCTION

Michael B. Gerhardt., M.D.

Wound Care:

Sutures remain on incision for one week Steri-strips remain on incision for two weeks Ace Bandage remains on your knee for two weeks Stocking remains on your knee for one week Ice at least four (4) times per day for thirty (30) minutes Shower with extremity covered for one week

You will be **partial weight bearing** for at least fourteen (14) days after surgery, using a cane or crutches.

Refrain from prolonged walking or standing for at least one (1) month post operatively

You will be **unable to drive a car** for at least three (3) days following surgery, and **unable to drive a car while on pain medication**

Exercise everyday following the instructions given to you at surgery (ie: straight leg raise, quad sets, knee bending, quad sets, Pedlar)

Medications: As prescribed for pain and inflammation

Physical Therapy Begins one week after surgery and lasts three (3) times a week for four (4) weeks

Post Op Visit Initial post op visit is one to three days following surgery